

¹CABIMER (Centro Andaluz de Biología Molecular y Medicina Regenerativa), Avda. Americo Vespucio s/n, Parque Científico y Tecnológico Cartuja, Sevilla, Spain; ²Stem Cells Therapies in Neurodegenerative Diseases Lab, Research Center "Principe Felipe", Valencia, Spain.; ³Spebo Medical, Leskovac, Serbia; ⁴Human Genetics, Faculty of Medical Sciences, University of Kragujevac, Serbia; ⁵Neuronal and Tissue Regeneration Lab, Research Center "Principe Felipe", Valencia, Spain; ⁶Institute of Experimental Medicine, AV CR, Prague, Czech Republic

*corresponding author: Slaven Erceg, Stem Cells Therapies in Neurodegenerative Diseases Lab, Research Center "Principe Felipe", C/ Eduardo Primo Yúfera 3, Valencia 46012, Spain, Tel. +34 963 289 680, Fax. +34 963 289 701, serceg@cipf.es

Received June 13, 2014; accepted for publication December 07, 2014; available online without subscription through the open access option.

©AlphaMed Press
1066-5099/2015/\$30.00/0

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/stem.1959

Reactive Astrocytes and Stem Cells in Spinal Cord Injury: Good Guys or Bad Guys?

DUNJA LUKOVIC¹, MIODRAG STOJKOVIC^{3,4}, VICTORIA MORENO-MANZANO⁵, PAVLA JENDELOVA⁶, EVA SYKOVA⁶, SHOMI S BHATTACHARYA¹ AND SLAVEN ERCEG^{1,2,6*}.

Key words. Glia • induced pluripotent stem cells • Neural differentiation • Neural stem cell • Stem cell transplantation • spinal cord injury

ABSTRACT

Spinal cord injury (SCI) usually results in long lasting locomotor and sensory neuron degeneration below the injury. Astrocytes normally play a decisive role in mechanical and metabolic support of neurons, but in the spinal cord they cause injury, exerting well-known detrimental effects that contribute to glial scar formation and inhibition of axon outgrowth. Cell transplantation is considered a promising approach for replacing damaged cells and promoting neuroprotective and neuroregenerative repair, but the effects of the grafted cells on local tissue and the regenerative properties of endogenous neural stem cells (eNSC) in the injured spinal cord are largely unknown. During the last two decades cumulative evidence from diverse animal models has indicated that reactive astrocytes in synergy with transplanted cells could be beneficial for injury in multiple ways, including neuroprotection and axonal growth. In this review we specifically focus on the dual opposing roles of reactive astrocytes in SCI and how they contribute to the creation of a permissive environment when combined with transplanted cells as the influential components for a local regenerative niche. Modulation of reactive astrocyte function might represent an extremely attractive new therapy to enhance the functional outcomes in patients. STEM CELLS 2014; 00:000–000

INTRODUCTION

Astrocytes are the most abundant glial cell type in the central nervous system (CNS), playing a decisive role in the mechanical and metabolic support of neurons, and

are involved primarily in regulating neural network activity, synaptic plasticity, blood flow, ionic and extracellular space volume homeostasis, and key neurotransmitter and ion concentrations[1, 2].

The definition of reactive astrocytes is related to radical change in the morphological and functional state of astrocytes as a response to a specific acute pathological condition such as spinal cord injury (SCI), brain injury, mechanical lesions of brain parenchyma, ischemia, and infection or neurodegenerative diseases[3]. In this review we analyze reactive astrocytes and their specific function in the spinal cord upon transplantation of pluripotent stem cells, trying to answer whether these cells exert a beneficial or detrimental function in regenerative processes after injury.

We specifically focus on the dual opposing role of reactive astrocytes in SCI and how they contribute to the creation of a permissive environment in synergy with transplanted cells as the key influential components in creating a local regenerative niche.

Reactive astrocytes in spinal cord injury: traditionally considered as bad guys

Spinal cord injury caused by mechanical trauma (contusion or compression) encompasses various processes that have been divided into two phases. The first phase is characterized by the induction of massive cell loss, with axonal, oligodendrocyte and neural cell death[4]. Massive loss of oligodendrocytes results in the inability of spared neurons to regenerate their axons[5], leading to neurological dysfunction. As a consequence of blood-brain barrier (BBB) damage and increased permeability, infiltrated cells from the blood invade the medullar tissue, triggering an inflammatory response. In the second phase, weeks after injury, different environmental cues associated with the inflammatory response and cell damage trigger existing astrocytes to undergo different morphological and molecular perturbations. These are mostly related to hypertrophic processes, increased cell proliferation and overexpression of glial fibrillary acidic protein (GFAP), vimentin and nestin[6], including the complex molecular, temporal and functional interconnections amongst the various signalling pathways (reviewed in [7],[3]).

These reactive astrocytes then interact with other cell types such as fibromeningeal cells and additional glial cells, contributing to the formation of a glial scar as a mechanical and chemical barrier to axonal regeneration. A more detailed description of glial scar formation can be found in the review by Silver and Miller[8]. The protein pattern of reactive astrocytes is also changed; reactive astrocytes overexpress GFAP and vimentin[6], the proteins that contribute to the formation of the glial scar[6], as well as S100 β , nestin, GLAST and GLT1 (Table 1). Furthermore, reactive astrocytes secrete inhibitory extracellular matrix molecules such as chondroitin sulfate proteoglycans (CSPGs), which inhibit axonal growth and regeneration[8], as well as tenascin, ephrin B2 and Slit proteins. These proteins secreted by reactive astrocytes in synergy with other cell types are traditionally considered to have detrimental roles by virtue of contributing to the formation of the physical and chemical barrier that impedes axonal regeneration. This negative

connotation of reactive astrocytes is an old concept and has been used to develop different strategies in order to regenerate the damaged tissue. One strategy is the manipulation of the glial scar using chondroitinase ABC to dissolve and reduce CSPG inhibition as a possible therapy for SCI[9], with moderate success. Application of chondroitinase ABC together with other intervention strategies such as peripheral nerve grafting[10], or Schwann cells and olfactory-ensheathing glia grafts[11] significantly increased axonal growth and improved locomotor outcomes, suggesting possible introduction of this approach in clinical practice[9].

Beneficial effects of reactive astrocytes: their neighbours make them good guys

The detrimental functions of reactive astrocytes mentioned previously are in fact responses to injury with the aim of protecting the nervous system and less affected tissue by healing the damaged area and halting further extension of injury and additional cell damage. At the same time, reactive astrocytes exert detrimental functions in the context of rendering the environment non-permissive for axonal repair and regeneration. *In vitro* and *in vivo* studies show that under normal conditions reactive astrocytes exert their protective role in the CNS in different ways such as: 1) active clearance of extracellular excitotoxic glutamate; 2) protection from ammonium toxicity; 3) protection from nitric oxide toxicity by producing glutathione; 4) degradation of amyloid β peptide; 5) stabilization of extracellular fluid and ion balance; 6) BBB repair; and 7) playing a role in energy (ATP) production via glucose transport. Upon SCI, besides expressing detrimental molecules, astrocytes contribute an important part in endogenous neuroprotection and also secrete growth-promoting neurotrophic factors. These are brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF), nerve growth factor (NGF), and basic fibroblast growth factor (FGF-2), as well as the extracellular matrix proteins laminin and fibronectin.

In the last 20 years, accumulating evidence from different deletion studies has revealed that reactive astrocytes exert a beneficial function upon injury, putting into question the concept of reactive astrocytes as a purely detrimental and negative phenomenon in SCI.

Genetically manipulated animal models provide further evidence that the loss or attenuation of reactive astrocytes worsens the outcome after CNS insults. While GFAP and/or Vimentin knockout (KO) mice show normal developmental and breeding patterns[12], they also exhibit low astroglial activity and increased plastic sprouting of supraspinal axons, including the reconstruction of circuits associated with functional outcomes in a hemisection model of SCI[13]. An elegant transgenic mouse model was developed by Sofroniew[14] that allowed the conditional ablation of GFAP-expressing cells using a herpes simplex virus type 1-thymidine kinase (suicide gene) under the control of the mouse GFAP promoter[14]. In this model it is possi-

ble to temporarily ablate dividing scar-forming reactive astrocytes using the antiviral drug ganciclovir in the context of different types of injury[14]. Such early ablation of reactive astrocytes limited scar formation in SCI models while the BBB was impaired, allowing increased immune cell infiltration, in particular macrophages, as well as neuronal death, demyelination, and impaired locomotor function[14]. The same effect of impaired recovery after SCI was observed in animals lacking matrix metalloproteinases (MMPs). MMPs proteolytic enzymes are involved in both, injury and repair mechanisms in the CNS and expressed in reactive astrocytes at the lesion border[15]. These results provide evidence that scar-forming astrocytes have a crucial role in neural protection and damage repair and are essential in tissue protection and preservation of function after mild or moderate SCI.

Okada et al. provided further evidence of a beneficial role of reactive astrocytes[16]. They applied a Cre-loxP system designed for conditional KO of signal transducer and activator of transcription 3 (STAT3) or cytokine signalling 3 suppressor (SOCS3), an intracellular signal transducer for various cytokines[16] involved in the up regulation of reactive astrocytes. The conditional ablation of STAT3 significantly decreased hypertrophy, overexpression of GFAP and the migration of reactive astrocytes, leading to increased inflammation, demyelination and neural disruption and poor locomotor outcome[16]. In addition, the deletion of SOCS3, which normally provides negative feedback on STAT3, displayed the opposite outcomes, reducing the lesion and improving motor symptoms after SCI[16]. Taken together, these findings reveal complex functions of reactive astrocytes that can be beneficial or detrimental for SCI depending on the signalling pathways involved. Proper modification of reactive astrocyte function after SCI could represent a new regeneration state.

The above described studies reveal the role of reactive astrocytes in preserving tissue integrity; however, little is known about how different cell replacement strategies modify the detrimental role of reactive astrocytes at the site of injury and contribute to improved recovery and enhanced axonal regeneration. We review cell transplantation approaches in tissue regeneration after SCI where the role of reactive astrocytes has been demonstrated. Schwann cells, foetal spinal cord, neural stem cells or neural derivatives of pluripotent stem cells are all considered promising sources for cell replacement therapy after SCI.

Schwann cells naturally populate the growth-permissive peripheral nervous system and their grafts seem to support axonal outgrowth by bridging the lesion gap following SCI, rendering the environment permissive for more extensive axonal outgrowth[17]. These early landmark transplantation studies showed successful regeneration of intraspinal, but not supraspinal axons. The latter have been regenerated when different combinatorial strategies were applied[18]. In most of these studies, analysis of the host spinal cord/Schwann

cells bridge interfaces revealed that astrocyte processes elongated in association with Schwann cells and regenerated axons, which correlated with locomotor outcome. For example, in the study of Guest et al.[19] the authors show that transplanted Schwann cells in combination with IN-1 anti-myelin antibodies induced migration of astrocytes into the site of a spinal cord transection and supported the axonal growth. Brainstem axons did not regenerate the lesion site and did not enter into the lesion if astrocytes failed to extend their processes. In contrast, Schwann cell grafts that exclude astrocytes are not invaded by CNS axons unless they are supplemented by exogenous neurotrophic factors. Elongation of reactive astrocytes in combination with Schwann cells can be triggered by glial-derived neurotrophic factor (GDNF) or by pre-treatment of glial-precursor astrocytes with bone-morphogenetic protein (BMP)[20]. This is a clear example of how appropriate stimulation or modification of the reactive astrocyte response after injury could be exploited for successful restoration of injured spinal cord.

After accumulating evidence that astrocytes are capable of responding to CNS injury by undergoing a process of reactive astrogliosis and providing support for cells and processes in the injured spinal cord, several studies have specifically exploited a cell transplantation strategy using astrocyte progenitors from embryonic or neonatal tissue in SCI models. These studies have addressed the phenotypic diversity or therapeutic application of glial restricted progenitors (GRP), particularly those derived from human foetal tissue. It has been shown that with proper *ex vivo* treatment these cells are capable of promoting neuroprotection and repairing damaged spinal cord[20, 21]. GRP can be isolated via the A2B5 surface marker from developing rat spinal cord[22], which are able to give rise to oligodendrocytes and astrocytes and exhibit self-renewal *in vitro*[22]. Upon treatment with either BMP or ciliary neurotrophic factor (CNTF), GRP gave rise to distinct types of astrocytes, but it was shown that only BMP-induced astrocytes exerted regenerative effects on axonal sprouting and locomotor recovery in the rat model of SCI[20, 23]. Human GRP transplanted into various types of SCI survived, migrated and differentiated toward astrocytes and oligodendrocytes[24]. In addition, these cells limited the formation of the glial scar through reducing CSPGs as well as enhancing axon sprouting [24] and promoted functional and sensory recovery[21]. Although GRPs represent a promising cell source for therapeutic application, the guidelines for their isolation, maintenance and differentiation remain to be standardized in order to optimize functional recovery after injury, which reveals the lack of well-established standards in preclinical studies using GRPs.

Early studies have shown that multipotent tissue grafts, such as foetal spinal cord[25], that were transplanted into damaged spinal cord, exerted neuroregenerative effects on the host tissue, by modifying scar formation in a manner associated with positive locomo-

tor outcome[26]. These studies highlight the possible contribution of reactive astrocytes to the regeneration of damaged spinal cord tissue upon transplantation of foetal grafts.

Neural stem cells (NSC) are a population of adult stem cells naturally present in adult neural tissue, primarily in the periventricular sub-ependymal layer and the subgranular zone of the dentate gyrus[27]. In the spinal cord, ependymal cells have neural stem cell properties and are located around the central canal[28]. Both brain and cord-derived NSC can be propagated *in vitro* as neurospheres and differentiate *in vitro* to form neurons, astrocytes and oligodendrocytes[29]. In response to injury, their proliferation capacity dramatically increases and constitutes the main pool of new glial cells in the injured spinal cord [30]. The regenerative effects of these cells after proper induction have been shown on different SCI models[29, 31]. In spite of their extensive capacity for neuronal differentiation *in vitro*, transplanted brain and spinal cord NSC *in vivo* differentiate almost exclusively into glia, with minimal neuronal differentiation observed[32]. In contrast, several transplantation studies of cells exhibiting astroglial hallmarks have been shown to promote functional recovery after SCI [33, 34].

We recently reported functional motor recovery after transplantation of spinal cord derived precursor cells, from spinal cord injured donors (ependymal stem-precursor cell induced by the injury, epSPCi), with induced cell proliferative capacity and glial preferential *in vivo* differentiation[29]. epSPCi in a contusion rat model showed a neuroprotective role by accompanying naked axons within the injured area, suggesting glial progenitor-derived neurotrophic support for axonal regeneration and cell survival[29]. Although NSC are considered to be ideal candidates for cell transplantation studies, due to their inherent neural commitment, low rates of tumorigenesis and the opportunity for autologous transplantation, their clinical application is questionable as a result of their limited availability and low proliferation rate.

Human pluripotent stem cells, encompassing human embryonic stem cells (hESC) and human induced pluripotent stem cells (hiPSC), hold great potential as a source for cell replacement therapies in humans. Human embryonic stem cells are pluripotent cells and can be derived from the inner cell mass of an early blastocyst[35]. A solution to the ethical concerns surrounding hESC was offered by the discovery of the Nobel prize winning technology of creating hiPSC from adult somatic cells by ectopic expression of a defined set of factors[36]. Human iPSC can be easily derived from individual patients allowing the development of customized stem cell therapies, generation of disease-specific stem cell lines and the possibility to correct the mutant gene in the target cells and return it to the patient[37]. Self-renewal and multi-lineage differentiation toward virtually any cell type in the human body are two unique properties that make hESC and iPSC the most

promising sources for tissue regeneration. hESC and iPSC are capable of differentiating toward all cell types, including neurons, glia, NSC, and motoneurons[38]. In the context of cell therapy for SCI, oligodendrocytes, astrocytes and neurons are of particular interest. As we already mentioned, SCI-induced massive cell death and loss of oligodendrocytes results in demyelination of spared axons leading to locomotor impairment. Delivery of early NPCs or oligodendrocyte progenitor cells (OPC) as a source for the remyelination processes, including the migration and mature differentiation of these cells, could be a promising strategy for spinal cord repair[39]. The regenerative properties of iPSC-derived neural progenitors in SCI[40-43] were found to take place through three possible mechanisms: remyelination, axonal regrowth and trophic support. Transplanted cells survive, migrate and differentiate toward all major neural cell lineages (astrocytes, neurons and oligodendrocytes)[44], corroborating the potential of hiPSC and hESC neural derivatives over adult stem cells[45]. Nevertheless, the mechanism of action of the transplanted cells is still unknown.

In the context of the beneficial role of reactive astrocytes in combination with pluripotent stem cell therapy, our recent observation of the regenerative potential of OPC when transplanted into a complete transection rat model of SCI, either alone or in combination with motoneuron progenitors (MP), is of particular interest[45]. We hypothesized that the potential of OPC and/or MP to rescue locomotor activity is due to the presence of heterogeneous cell types present in transplanted progenitors[46]. We identify their possible mechanism through the modification of inhibitory properties of reactive astrocytes[46]. In this study we analysed the microenvironment of the lesion site, revealing the molecules and processes that could point to the possible mechanism of action. We showed enhanced astrogliosis four months after transplantation of OPC and MP in the acute phase of SCI, possibly as a result of Notch and JAK/STAT signalling activation. At that time point, the GFAP-negative area in the epicentre of the injury site was significantly reduced in transplanted animals, suggesting that JAK/STAT and Notch signalling enhanced astrogliosis in the lesion site resulting in lesion compaction, as has been previously observed[47]. In transplanted rats we found improved growth of serotonergic and dopaminergic axons in close proximity with transplanted cells and GFAP⁺ astrocytes. It seems that the prompt contraction of the lesion favours hESC-derived neuronal survival and differentiation and correlates with improved locomotor recovery in these animals[16]. This study corroborates the role of astrogliosis in spinal cord regeneration[14, 23], suggesting that neural protective factors secreted from transplanted cells in synergy with reactive astrocytes could create the neuroprotective environment for neurogenesis observed in transplanted cells[45]. This particular niche created by transplanted cells seems to alter the detrimental properties of reactive astrocytes. The increase in the expres-

sion of beneficial molecules such as NGF, laminins, fibronectin and neurotrophins, and decreases in detrimental genes such as CSPG, TENASCINS and genes included in SLIT/ROBO signalling in the lesion site resulted in less inhibitory reactive astrocytes, making them permissive to axonal growth, neuronal progenitor survival and differentiation[46]. This in turn is associated with the presence of human neurons in the lesion site[45]. This finding corroborates reports of iPS progenitor-derived astrocyte promotion of axonal regrowth via a growth-permissive substrate[43], which is involved in specifically creating a permissive environment where beneficial molecules such as BDNF are expressed. This was recently demonstrated in a study by Romanyuk et al.[41]. The absence of a growth-promoting cell population in the control group resulted in a lack of axonal regeneration.

Finally, growing lines of evidence suggest that astroglia lineages host neural stem-like cells[48] revealing their *in vitro* capacity to self-renew and give rise to progenies including all three neural lineages[7, 48] when grown as neurospheres. Recently, it has been shown that injury reactivates stem cell potential of reactive astrocytes[49] through sonic hedgehog (SHH) protein[50]. It is of particular interest to understand the mechanisms behind the proliferative response of reactive glia in different injury settings in order to assess the ability of these NSC-like cells to provide endogenous repair for therapeutic purposes.

Therapeutic interventions to make reactive astrocytes good guys

As soon as reactive astrocytes had been recognized as key regulators of spinal cord injury, specific molecular aspects of reactive astrogliosis were targeted for the purpose of therapeutic manipulations[51]. It has been identified that epigenetic regulation of gene expression can be of great therapeutic importance. Acetylation and deacetylation of histones using histone deacetylases (HDACs) and histone acetyltransferases (HATs), respectively, affecting the global gene expression pattern in cells, can be of special interest in the neurological field. For example, pharmacological inhibition of HDAC with valproic acid (VA) increases histone acetylation and improves pathological conditions in various neurological diseases, including SCI[52]. VA was shown to stimulate the release of neurotrophic factors from astrocytes [53], at the same time decreasing autophagy and reducing the expression of inflammatory mediators including tumor necrosis factor- α [54]. Curcumin, a major curcumanoid in the spice turmeric and HAT inhibitor is also a prospective drug for neurological diseases. The effects of curcumin have been generally described as anti-inflammatory, anti-apoptotic and anti-oxidative, improving neurological function in the injured spinal cord of rats[55]. Curcumin exerts its function through inhibiting astrocyte reactivation, which may be beneficial for neuronal survival, as claimed in a study by Lin et al.[55] and other study of various types of injuries[56].

However future studies are needed to clarify the roles of curcumin in the regulation of signalling pathways and gene expression profiles in astrocytes.

Other potential targets for astrocyte functional modification include Aquaporin 4, a water channel responsible for bi-directional, blood-brain water transport, normally localized on astrocyte endfeet, or different connexins or potassium channels localized to the same site, responsible for maintaining the resting membrane potential and needed for proper K^+ buffering. In the context of astrocyte function, these targets have been well reviewed by Hamby et al.[51].

A recent article by Renault-Mihara et al.[57] showed that the beneficial role of reactive astrocytes can be triggered in SCI by inhibiting Glycogen synthase kinase 3 (GSK-3), a serine/threonine protein kinase, resulting in improved locomotor activity in mice. This study clearly demonstrates the stimulation of astrocyte migration as a feasible therapeutic strategy for traumatic injury in the central nervous system.

CONCLUSION

The growing evidence summarized here indicates that reactive astrocytes have both active and passive roles in regenerative processes after SCI. Reactive astrocytes *per se* cannot repair damaged SCI, but can contribute to healing the area by protecting the lesion site from any further extension of damage. It seems that only in the proper environment can these cells modify their own properties and create a permissive niche for endogenous events and exogenous interventions to regenerate the damaged tissue. Also in this review, we reported the novel function of transplanted hESC derivatives on modifying properties of reactive astrogliosis, which was classically considered as detrimental to axonal regeneration, but has also been shown to contribute to a favourable environment for neuronal differentiation of transplanted cells. Overall, a future repair strategy based on combined cell and drug therapy with pluripotent stem cell-derived neural cells that minimize the inhibitory properties of astrocytes while maximizing their growth-promoting properties would be extremely attractive for the treatment of SCI. It will be challenging to 'fine tune' the modification of reactive astrocyte function via a selective blockade of molecules that inhibit axonal outgrowth while allowing the formation of a physical barrier to protect the intact tissue.

ACKNOWLEDGMENTS

This work was supported by funds for research from "Miguel Servet" contract of Instituto de Salud Carlos III of Spanish Ministry of Science and Innovation (SE), Fund for Health of Spain PI10-01683 (VM), Junta de Andalucía PI-0113-2010 (SE) and Wings for Life Foundation (SE), Czech National Foundation GA CR P304/12/G069 (PJ) and LO1309 of the MEYS of the CR under the NPU I pro-

gramme from the European Regional Development Fund (ES).

AUTHOR CONTRIBUTIONS

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

The authors indicate no potential conflicts of interest.

D.L.: manuscript writing, final approval of manuscript; V.M.-M.: Final approval of manuscript; S.E.: Conception and design, manuscript writing, final approval of manuscript; M.S.: Final approval of manuscript; S.S.B.: Final approval of manuscript; P.J.: Final approval of manuscript; E.S.: Final approval of manuscript

REFERENCES

- Ransom B, Behar T, Nedergaard M. New roles for astrocytes (stars at last). *Trends Neurosci.* 2003;26:520-522.
- Sykova E, Nicholson C. Diffusion in brain extracellular space. *Physiological reviews.* 2008;88:1277-1340.
- Eddleston M, Mucke L. Molecular profile of reactive astrocytes—implications for their role in neurologic disease. *Neuroscience.* 1993;54:15-36.
- Grossman SD, Rosenberg LJ, Wrathall JR. Temporal-spatial pattern of acute neuronal and glial loss after spinal cord contusion. *Exp Neurol.* 2001;168:273-282.
- Kakulas BA. The applied neuropathology of human spinal cord injury. *Spinal Cord.* 1999;37:79-88.
- Pekny M, Nilsson M. Astrocyte activation and reactive gliosis. *Glia.* 2005;50:427-434.
- Tucker BA, Scheetz TE, Mullins RF et al. Exome sequencing and analysis of induced pluripotent stem cells identify the cilia-related gene *male germ cell-associated kinase (MAK)* as a cause of retinitis pigmentosa. *Proc Natl Acad Sci U S A.* 2011;108:E569-576.
- Silver J, Miller JH. Regeneration beyond the glial scar. *Nat Rev Neurosci.* 2004;5:146-156.
- Bradbury EJ, Carter LM. Manipulating the glial scar: chondroitinase ABC as a therapy for spinal cord injury. *Brain research bulletin.* 2011;84:306-316.
- Yick LW, Wu W, So KF et al. Chondroitinase ABC promotes axonal regeneration of Clarke's neurons after spinal cord injury. *Neuroreport.* 2000;11:1063-1067.
- Fouad K, Schnell L, Bunge MB et al. Combining Schwann cell bridges and olfactory-ensheathing glia grafts with chondroitinase promotes locomotor recovery after complete transection of the spinal cord. *J Neurosci.* 2005;25:1169-1178.
- Pekny M, Wilhelmsson U, Bogestal YR et al. The role of astrocytes and complement system in neural plasticity. *International review of neurobiology.* 2007;82:95-111.
- Menet V, Prieto M, Privat A et al. Axonal plasticity and functional recovery after spinal cord injury in mice deficient in both glial fibrillary acidic protein and vimentin genes. *Proc Natl Acad Sci U S A.* 2003;100:8999-9004.
- Faulkner JR, Herrmann JE, Woo MJ et al. Reactive astrocytes protect tissue and preserve function after spinal cord injury. *J Neurosci.* 2004;24:2143-2155.
- Hsu JY, McKeon R, Goussev S et al. Matrix metalloproteinase-2 facilitates wound healing events that promote functional recovery after spinal cord injury. *The Journal of neuroscience : the official journal of the Society for Neuroscience.* 2006;26:9841-9850.
- Okada S, Nakamura M, Katoh H et al. Conditional ablation of Stat3 or Socs3 discloses a dual role for reactive astrocytes after spinal cord injury. *Nat Med.* 2006;12:829-834.
- Campbell G, Hutchins K, Winterbottom J et al. Upregulation of activating transcription factor 3 (ATF3) by intrinsic CNS neurons regenerating axons into peripheral nerve grafts. *Exp Neurol.* 2005;192:340-347.
- Xu XM, Guenard V, Kleitman N et al. A combination of BDNF and NT-3 promotes supraspinal axonal regeneration into Schwann cell grafts in adult rat thoracic spinal cord. *Exp Neurol.* 1995;134:261-272.
- Guest JD, Rao A, Olson L et al. The ability of human Schwann cell grafts to promote regeneration in the transected nude rat spinal cord. *Exp Neurol.* 1997;148:502-522.
- Davies JE, Proschel C, Zhang N et al. Transplanted astrocytes derived from BMP- or CNTF-treated glial-restricted precursors have opposite effects on recovery and allodynia after spinal cord injury. *Journal of biology.* 2008;7:24.
- Alexanian AR, Svendsen CN, Crowe MJ et al. Transplantation of human glial-restricted neural precursors into injured spinal cord promotes functional and sensory recovery without causing allodynia. *Cytotherapy.* 2011;13:61-68.
- Dietrich J, Noble M, Mayer-Proschel M. Characterization of A2B5+ glial precursor cells from cryopreserved human fetal brain progenitor cells. *Glia.* 2002;40:65-77.
- Davies JE, Huang C, Proschel C et al. Astrocytes derived from glial-restricted precursors promote spinal cord repair. *Journal of biology.* 2006;5:7.
- Hill CE, Proschel C, Noble M et al. Acute transplantation of glial-restricted precursor cells into spinal cord contusion injuries: survival, differentiation, and effects on lesion environment and axonal regeneration. *Exp Neurol.* 2004;190:289-310.
- Bernstein JJ, Goldberg WJ. Graft derived reafferentation of host spinal cord is not necessary for amelioration of lesion-induced deficits: possible role of migrating grafted astrocytes. *Brain research bulletin.* 1989;22:139-146.
- Houle JD, Reier PJ. Transplantation of fetal spinal cord tissue into the chronically injured adult rat spinal cord. *J Comp Neurol.* 1988;269:535-547.
- Weiss S, Dunne C, Hewson J et al. Multipotent CNS stem cells are present in the adult mammalian spinal cord and ventricular neuroaxis. *J Neurosci.* 1996;16:7599-7609.
- Hawrylyk GW, Fehlings MG. The center of the spinal cord may be central to its repair. *Cell Stem Cell.* 2008;3:230-232.
- Moreno-Manzano V, Rodriguez-Jimenez FJ, Garcia-Rosello M et al. Activated spinal cord ependymal stem cells rescue neurological function. *Stem Cells.* 2009;27:733-743.
- Barnabe-Heider F, Goritz C, Sabelstrom H et al. Origin of new glial cells in intact and injured adult spinal cord. *Cell Stem Cell.* 2010;7:470-482.
- Enzmann GU, Benton RL, Talbott JF et al. Functional considerations of stem cell transplantation therapy for spinal cord repair. *J Neurotrauma.* 2006;23:479-495.
- Cao Q, Benton RL, Whitemore SR. Stem cell repair of central nervous system injury. *J Neurosci Res.* 2002;68:501-510.
- Cao Q, Xu XM, Devries WH et al. Functional recovery in traumatic spinal cord injury after transplantation of multilineurotrophin-expressing glial-restricted precursor cells. *J Neurosci.* 2005;25:6947-6957.
- Hasegawa K, Chang YW, Li H et al. Embryonic radial glia bridge spinal cord lesions and promote functional recovery following spinal cord injury. *Exp Neurol.* 2005;193:394-410.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS et al. Embryonic stem cell lines derived from human blastocysts. *Science.* 1998;282:1145-1147.
- Takahashi K, Tanabe K, Ohnuki M et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell.* 2007;131:861-872.
- Raya A, Rodriguez-Piza I, Guenechea G et al. Disease-corrected haematopoietic progenitors from Fanconi anaemia induced pluripotent stem cells. *Nature.* 2009;460:53-59.
- Erceg S, Ronaghi M, Stojkovic M. Human embryonic stem cell differentiation toward regional specific neural precursors. *Stem Cells.* 2009;27:78-87.
- Keirstead HS, Nistor G, Bernal G et al. Human embryonic stem cell-derived oligodendrocyte progenitor cell transplants remyelinate and restore locomotion after spinal cord injury. *J Neurosci.* 2005;25:4694-4705.
- Fujimoto Y, Abematsu M, Falk A et al. Treatment of a mouse model of spinal cord injury by transplantation of human induced pluripotent stem cell-derived long-term self-

- renewing neuroepithelial-like stem cells. **Stem Cells**. 2012;30:1163-1173.
- 41 Romanyuk N, Amemori T, Turnovcova K et al. Beneficial effect of human induced pluripotent stem cell-derived neural precursors in spinal cord injury repair. **Cell transplantation**. 2014;Epub ahead of print;DOI: 10.3727/096368914X096684042.
- 42 Tsuji O, Miura K, Fujiyoshi K et al. Cell therapy for spinal cord injury by neural stem/progenitor cells derived from iPS/ES cells. **Neurotherapeutics**. 2011;8:668-676.
- 43 Tsuji O, Miura K, Okada Y et al. Therapeutic potential of appropriately evaluated safe-induced pluripotent stem cells for spinal cord injury. **Proc Natl Acad Sci U S A**. 2010;107:12704-12709.
- 44 Nori S, Okada Y, Yasuda A et al. Grafted human-induced pluripotent stem-cell-derived neurospheres promote motor functional recovery after spinal cord injury in mice. **Proc Natl Acad Sci U S A**. 108:16825-16830.
- 45 Erceg S, Ronaghi M, Oria M et al. Transplanted oligodendrocytes and motoneuron progenitors generated from human embryonic stem cells promote locomotor recovery after spinal cord transection. **Stem Cells**. 2010;28:1541-1549.
- 46 Lukovic D, Valdes-Sanchez L, Sanchez-Vera I et al. Brief Report: Astroglial Promotes Functional Recovery of Completely Transected Spinal Cord Following Transplantation of hESC-Derived Oligodendrocyte and Motoneuron Progenitors. **Stem Cells**. 2014;32:594-599.
- 47 Tsuda M, Kohro Y, Yano T et al. JAK-STAT3 pathway regulates spinal astrocyte proliferation and neuropathic pain maintenance in rats. **Brain**. 2011;134:1127-1139.
- 48 Riazifar H, Jia Y, Chen J et al. Chemically induced specification of retinal ganglion cells from human embryonic and induced pluripotent stem cells. **Stem cells translational medicine**. 2014;3:424-432.
- 49 Riesenberger AN, Le TT, Willardson MJ et al. Pax6 regulation of Math5 during mouse retinal neurogenesis. **Genesis**. 2009;47:175-187.
- 50 Le TT, Wroblewski E, Patel S et al. Math5 is required for both early retinal neuron differentiation and cell cycle progression. **Dev Biol**. 2006;295:764-778.
- 51 Hamby ME, Sofroniew MV. Reactive astrocytes as therapeutic targets for CNS disorders. **Neurotherapeutics**. 2010;7:494-506.
- 52 Hao HH, Wang L, Guo ZJ et al. Valproic acid reduces autophagy and promotes functional recovery after spinal cord injury in rats. **Neuroscience bulletin**. 2013.
- 53 Chen PS, Peng GS, Li G et al. Valproate protects dopaminergic neurons in midbrain neuron/glia cultures by stimulating the release of neurotrophic factors from astrocytes. **Molecular psychiatry**. 2006;11:1116-1125.
- 54 Lee JY, Kim HS, Choi HY et al. Valproic acid attenuates blood-spinal cord barrier disruption by inhibiting matrix metalloproteinase-9 activity and improves functional recovery after spinal cord injury. **J Neurochem**. 2012;121:818-829.
- 55 Lin MS, Lee YH, Chiu WT et al. Curcumin provides neuroprotection after spinal cord injury. **The Journal of surgical research**. 2011;166:280-289.
- 56 Laird MD, Sukumari-Ramesh S, Swift AE et al. Curcumin attenuates cerebral edema following traumatic brain injury in mice: a possible role for aquaporin-4? **J Neurochem**. 2010;113:637-648.
- 57 Renault-Mihara F, Katoh H, Ikegami T et al. Beneficial compaction of spinal cord lesion by migrating astrocytes through glycogen synthase kinase-3 inhibition. **EMBO molecular medicine**. 2011;3:682-696.

Figure 1. Schematic overview of astrocyte modulation upon cell transplantation. Spinal cord injury triggers the migration of reactive astrocytes (1). Stem cell progenitors transplanted during the acute phase of injury differentiate toward neurons (2), which reconnect the damaged tissue, and oligodendrocytes (3) that remyelinate new neurons and sprouted axons (4). Transplanted cells modulate the detrimental role of reactive astrocytes and create a permissive environment for the upregulation of beneficial proteins such as laminins (L) and fibronectins (F).




Table 1. Outline of markers of proteoplasmatic astrocytes and reactive ones. GFAP- Glial fibrillary acidic protein; CSPGs- Chondroitin sulfate proteoglycans; GLAST- Glutamate-aspartate transporter; GLT1- Glial Glutamate Transporter 1; BLBP- lipid-binding protein; AQP4- Aquaporin 4; ↑Overexpression.

Proteoplasmatic astrocytes	Reactive astrocytes
GFAP	GFAP ↑
S100β	S100β ↑
GLAST	GLAST ↑
GLT1	GLT1 ↑
BLBP	BLBP ↑
AQP4	AQP4↑
	CSPGs↑
	Ephrins↑
	Chemokines↑
	Vimentin↑
	Nestin↑